

# Identification of a cellulose synthase-associated protein required for cellulose biosynthesis

Ying Gu<sup>a,1</sup>, Nick Kaplinsky<sup>b</sup>, Martin Bringmann<sup>c</sup>, Alex Cobb<sup>d</sup>, Andrew Carroll<sup>a,e</sup>, Arun Sampathkumar<sup>c</sup>, Tobias I. Baskin<sup>d</sup>, Staffan Persson<sup>c</sup>, and Chris R. Somerville<sup>a,2</sup>

<sup>a</sup>Energy Biosciences Institute, University of California, Berkeley, CA 94720; <sup>b</sup>Department of Biology, Swarthmore College, Swarthmore, PA 19081; <sup>c</sup>Max-Planck-Institute for Molecular Plant Physiology, 14476 Potsdam, Germany; <sup>d</sup>Biology Department, University of Massachusetts, Amherst, MA 01003; and <sup>e</sup>Department of Biology, Stanford University, Stanford, CA 94305

Contributed by Chris R. Somerville, May 24, 2010 (sent for review May 8, 2010)

**Cellulose synthase-interactive protein 1 (CS11) was identified in a two-hybrid screen for proteins that interact with cellulose synthase (CESA) isoforms involved in primary plant cell wall synthesis. CS11 encodes a 2,150-amino acid protein that contains 10 predicted Armadillo repeats and a C2 domain. Mutations in CS11 cause defective cell elongation in hypocotyls and roots and reduce cellulose content. CS11 is associated with CESA complexes, and *csi1* mutants affect the distribution and movement of CESA complexes in the plasma membrane.**

cell walls | CESA complexes | cell expansion | microtubules | polysaccharides

Cellulose is synthesized at the plasma membrane by hexameric protein complexes with a diameter of 25–30 nm when observed by freeze-fracture electron microscopy in algae, moss, and vascular plants (1–4). The only known component of the complexes is cellulose synthase (CESA), which is represented by 10 isoforms in *Arabidopsis* (5). Genetic studies indicate at least three *Arabidopsis* CESA isoforms are required for primary cell wall synthesis (6, 7). Lesions in CESA1 (*rsw1*), CESA3 (*cev1*), or CESA6 (*prc1*) lead to a deficiency in elongation in dark-grown seedlings (8–10). Genetic and biochemical studies demonstrating interactions between CESAs led to a heteromeric model of cellulose synthesis (11, 12) in which the complexes are composed of at least three functionally nonredundant CESA isoforms. It generally is accepted that in *Arabidopsis* CESA1, CESA3, and CESA6 or CESA6-like proteins are required for functional primary cell wall complexes, whereas CESA4, CESA7, and CESA8 are required for functional secondary cell wall complexes. However, the exact number of CESA proteins contained within the complex, their stoichiometry, and their specific interactions are unknown, and no other components of the complex have been reported.

Recent advances in cell biology and microscopy allow imaging of CESA complexes in live tissues. At least two of the three primary CESAs (CESA3 and CESA6) are functional when labeled with GFP and its derivatives. Both GFP-CESA3 and YFP-CESA6 were observed at the plasma membrane as discreet particles that move along linear trajectories coincident with underlying cortical microtubules (7, 13). CESA particles move bidirectionally with an average velocity of about 350 nm/min corresponding to the addition of ~700 glucose residues per glucan chain per minute (13). CESA particle dynamics are sensitive to osmotic stress and to several drugs that affect cytoskeleton and cellulose synthesis. The observation that perturbation of microtubule polymerization by oryzalin affects the overall distribution and motility of CESA particles supports models in which the microtubules guide the deposition of cellulose. However, CESA particles appear to have an intrinsic level of organization that is evident when microtubules are completely depleted (13).

Genetic screens for mutants deficient in cellulose have implicated a number of proteins in the overall process of cellulose biosynthesis. Mutations in *KORRIGAN* (*KOR*), which encodes an endo- $\beta$ -1,4-glucanase, exhibit deficiencies in cell elongation and reduced cellulose production (14). *KOR*-like proteins from

*Brassica napus* and poplar exhibit cellulase activity in vitro. However, the exact role of cellulase in cellulose synthesis is unknown. Additional cellulose-deficient mutants include *cobra*, *kobito*, *pom1*, *rsw3*, *fragile fiber1*, and *fragile fiber2*, none of which has been assigned a clear mechanistic function in cellulose synthesis. Here we report the identification of a protein involved in cellulose synthesis that appears to be associated with primary CESA complexes. Identification of this protein opens an avenue in ongoing efforts to understand the mechanism of cellulose synthesis.

## Results

**Cellulose Synthase-Interactive Protein 1 Interacts with Multiple Primary CESAs.** To explore whether additional proteins may be required for cellulose biosynthesis, we performed yeast two-hybrid screens to identify proteins that physically interact with CESA1, -3, and -6. Using 541 amino acids of the putative catalytic domain of CESA6 as bait, we identified a protein referred to as “cellulose synthase-interactive protein 1” (CS11; At2g22125) (Fig. 1A). To confirm the interaction of CESA6 and CS11, we subcloned the prey into a GAL4 activation domain (GAL4-AD) and fused the putative catalytic domain of CESA6 with GAL4 binding domain (GAL4-BD). Coexpression of these constructs in yeast resulted in the appearance of  $\beta$ -galactosidase (GUS) activity, confirming the interaction between CESA6 and CS11. CESA1 and CESA3 also showed positive interactions with CS11, although the CESA3 interaction appeared to be weaker than the CESA1 or CESA6 interaction (Fig. 1B).

*CS11* was identified previously as one of the genes that is coregulated transcriptionally with the primary CESAs (15). In addition to *CS11*, several genes that affect cellulose deposition, including *COBRA* and *CTL1/POM1*, also are coexpressed with the *CESA* genes (Fig. 1C). Consistent with coexpression analysis, transgenic plants in which *GUS* was placed under the control of the 1.4-kb promoter region upstream of the *CS11* gene exhibited a pattern of GUS activity (Fig. 1D, F, and H and Fig. S1) similar to that seen with promoter:GUS fusions of primary wall *CESA* genes (Fig. 1E, G, and I). In addition, the *CS11* promoter drives GUS expression in floral tissues, rosette leaves, roots, and pollen (Fig. S1), indicating that the *CS11* expression pattern is similar to that of CESAs throughout development.

Author contributions: Y.G., N.K., M.B., A. Cobb, A. Carroll, T.I.B., S.P., and C.R.S. designed research; Y.G., N.K., M.B., A. Cobb, A. Carroll, and A.S. performed research; A. Cobb contributed new reagents/analytic tools; Y.G., N.K., M.B., A. Cobb, A. Carroll, A.S., T.I.B., S.P., and C.R.S. analyzed data; and Y.G., N.K., M.B., A. Cobb, A. Carroll, A.S., T.I.B., S.P., and C.R.S. wrote the paper.

The authors declare no conflict of interest.


<sup>1</sup>Present address: Department of Biochemistry and Molecular Biology, Pennsylvania State University, University Park, PA 16802.

<sup>2</sup>To whom correspondence should be addressed. E-mail: crs@berkeley.edu.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1007092107/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1007092107/-DCSupplemental).








**Fig. 2.** CSI1 is required for normal cell expansion. (A) Schematic representation of six T-DNA insertion sites in *csi1*. Exons are represented by black lines, and introns are shown by breaks. (B) Morphology of 4-d-old dark-grown seedlings: (Left to Right) Col-0 (wild-type) and *csi1-1*, *csi1-2*, *csi1-3*, *csi1-4*, *csi1-5*, and *csi1-6* mutants. (Scale bar: 2 mm.) (C and D) Hypocotyl length (C) and growth rate (D) of dark-grown wild-type (Col-0) plants and *csi1-3*, *csi1-6*, and *prc1-1* mutants. Data were collected from the measurement of ~50 seedlings for each genotype. Error bars represent SE (absent error bars were obscured by symbols). (E) SEM of dark-grown hypocotyls in wild-type plants and *csi1* mutants: (Left to Right) *Arabidopsis thaliana* Columbia (Col-0), *csi1-3*, and *csi1-6* mutants. Colors outline one epidermal cell. (Scale bar: 100 μm.) (F) Cellulose content was reduced in *csi1* mutants.  $n = 5$ . Error bars represent SE.

fertility of *csi1* mutants was caused mainly by pollen defects (Table S1). Therefore, we examined the surface structure of the mature pollen grains by SEM. Although all wild-type pollen grains displayed typical morphology, very few typical pollen grains were observed in the *csi1* alleles. Instead, the majority of the *csi1* pollen grains displayed irregular or collapsed cell wall morphologies (Fig. S3 I and K). Similar observations also have been reported for null alleles of *CESA1* and *CESA3* (6).


**CSI1 Is Associated with CESA Complexes.** To investigate the subcellular localization of CSI1, we generated both N- and C-terminal translational fusions of red fluorescent protein (RFP) to the CSI1 protein under the native *CSII* promoter in the *csi1-6*-mutant background. Both N- and C-terminal constructs complemented the mutant phenotypes of reduced hypocotyl elongation and reduced expansion anisotropy, indicating that the fusion protein was functional (Fig S4). Observation of epidermal cells in dark-grown hypocotyls by spinning disk confocal microscopy revealed that RFP-CSI1 was detected as distinct particles at the plasma membrane (Fig. 3A and Movie S1). Similar to YFP-CESA6, RFP-CSI1 particles were organized into linear arrays. To assess whether RFP-CSI1 particles move bidirectionally like those of YFP-CESA6, directional bias was analyzed by calculating the dot product of each particle relative to the dominant axis of potential particle bias and correlating this dot product with the velocity of each particle (Fig. 3E). Greater differences between particle velocities going with and against the major axis result in a larger slope in a linear regression of the plot. As seen in Fig. 3E, RFP-CSI1 particles ( $n = 917$ ) travel bidirectionally with no bias relative to the major axis. The average



**Fig. 3.** CSI1 is localized to CESA-like particles in dark-grown hypocotyls cells. (A–D) Optical sections of epidermal cells in 3-d-old dark-grown hypocotyls expressing RFP-CSI1 (A and C) and YFP-CESA6 (B and D). Focal planes of the plasma membrane (A and B) and lower cortex (C and D) are shown. (Scale bar: 5 μm.) (E) Plot of RFP-CSI1 particle velocity vs. its dot product (i.e., scalar product; see Methods) with the direction of bias. Red line indicates the linear regression of the plot. A slope of 0 indicates no bias in direction. (F) Histogram of measured RFP-CSI1 particle velocities. The mean is 416 nm/min ( $n = 917$ ). (G–I) Localization of GFP-CESA3 (G), RFP-CSI1 (H), and merge (I). (Scale bar: 5 μm.)

velocity of RFP-CSI1 particles in epidermal cells in dark-grown hypocotyls was 416 nm/min (range 63–860 nm/min; Fig. 3F). The velocity of RFP-CSI1 is similar to that reported for YFP-CESA6 (13). Unlike YFP-CESA6, which is associated with the Golgi complex in addition to the plasma membrane (Fig. 3B and D) (13), RFP-CSI1 was not detected in the cytoplasmic compartments (Fig. 3C). In plants containing both GFP-CESA3 and RFP-CSI1, the RFP-CSI1 particles were substantially colocalized with GFP-CESA3 (Fig. 3G–I).

**Lesions in *csi1* Affect the Distribution and Motility of YFP-CESA6.** To monitor the dynamics of CESA complexes in *csi1* mutants directly, we introduced a homozygous *csi1-3* allele into a YFP-CESA6 line (13). Using the YFP-CESA6 marker, CESA complexes can be observed at the plasma membrane as distinct punctuate particles that move at constant rates along linear tracks (13). Although the CESA particles in the plasma membranes of wild-type (control) epidermal cells were organized into linear arrays, the distribution of the CESA particles appeared to be disorganized in plasma membranes of the *csi1-3* background (Fig. 4C and Movie S2). In contrast, CESA particles in the internal cell layers were not affected in *csi1-3* mutants. We did not observe any significant differences in



**Fig. 4.** YFP-CESA6 dynamics are altered in *csil-3* mutants. YFP-CESA6 localization in dark-grown hypocotyls cells is shown in wild-type plants (A and B) and *csil-3* mutants (C and D). (A) A single optical section acquired at the plane of plasma membrane in wild-type plants. (B) Average of 61 frames (duration: 2 min; 2-s interval) showing movement of labeled particles along linear trajectories. (C) Single optical section of YFP-CESA6 in *csil-3* mutant. (D) Average of 61 frames (duration: 2 min; 2-s interval) in *csil-3* mutant. (Scale bars: 5  $\mu$ m.) (E) Histogram of measured particle velocities. The mean velocity is 365 nm/min in control plants ( $n = 318$ ) and 132 nm/min in *csil-3* mutants ( $n = 225$ ).

the distribution or motility of CESA particles associated with the Golgi complex in *csil-3* mutants compared with control lines (Golgi in the *csil-3* mutants moved at  $6021 \pm 405$  nm/min vs.  $5214 \pm 982$  nm/min in wild-type plants). In control cells, CESA particles associated with the plasma membrane migrated with an average velocity of  $365 \pm 45$  nm/min ( $n = 318$ ). In *csil-3* mutants, however, the average velocity of CESA particles was reduced to  $132 \pm 52$  nm/min ( $n = 225$ ; Fig. 4E). These observations are reflected clearly in time-averaged projections of the CESA particles (Fig. 4 B and D). In control cells, motile particles form linear trajectories along the axes of the particle arrays. Although CESA particles still move bidirectionally in *csil-3* mutants (as assessed by Imaris software), linear tracks in time-averaged images were much shorter.

**Uniformity of Cellulose Microfibrils Is Affected in *csil* Mutants.** To investigate effects of the *csi* mutations on the arrangement of cellulose microfibrils, we observed the longitudinal cell walls of wild-type and *csil-1* roots using circularly polarized light enhanced with the universal compensator (26). Polarized light can be used to assay the abundance and orientation of cellulose microfibrils because of their partial crystallinity. Overall, the cell walls appeared similar in both genotypes (Fig. 5 A and B). To determine whether there was a quantitative difference in microfibril abundance or orientation, we quantified the retardance of the cell walls and the azimuth of the crystalline elements. The average retardance was about 1 nm, similar to that reported previously for *Arabidopsis* cell walls (27), and there was no significant difference between the genotypes in the three delineated zones of the root (Fig. 5C). Similarly, in the three root zones of

both genotypes, the azimuths averaged to  $90^\circ$  transverse to the long axis of the root. However, for the elongation and mature zones, the variability among azimuth measurements made in individual roots was significantly greater in *csil-1* mutants than in wild-type roots, suggesting that microfibril alignment in these cell walls had become less uniform.

## Discussion

The CSI1 protein is the first non-CESA protein associated with CESA complexes. Several lines of evidence led us to hypothesize that CSI1 exerts a direct effect on cellulose synthesis through its association with CESA complexes. First, CSI1 physically interacts with multiple primary CESAs in yeast two-hybrid assays, and CSI1 is transcriptionally coregulated with several of the *CESA* genes involved in primary cell wall synthesis, but there was no obvious association with CESAs involved in secondary cell wall synthesis. Additionally, CSI1 colocalizes with primary CESA complexes, and *csil* mutations affect the distribution and movement of CESA complexes, resulting in strongly reduced rates of CESA complex movement. If we assume that the length of cellulose microfibrils is affected by the velocity and lifetimes of CESA particles, the cellulose deficiency and the associated swelling phenotype can be attributed to the effect of the *csil* mutations on the activity of the CESA complexes. Additionally, the *csil* mutations appeared to decrease the degree to which cellulose microfibrils are coaligned. Polarized light is sensitive to the azimuth at which the optical axis of the crystalline sample is oriented. In both *csil* mutants and wild-type roots the average azimuth of cellulose microfibrils was transverse to the long axis of the root. However, the SD of the azimuth measurements was larger in *csil* mutants than in wild-type roots. In other words, microfibril alignment, on a scale greater than that of a wavelength, is noisier in *csil* mutants than in wild-type roots. A similar reduction in the uniformity of microfibril alignment across the root has been reported for treatment with low concentrations of the microtubule inhibitor oryzalin (27). The decrease in cellulose organization in *csil* mutants indicates that CSI1 may participate in the mechanisms responsible for organizing the deployment of cellulose microfibrils in primary walls.


CSI1 belongs to a family of highly conserved land plant-specific proteins. CSI1 contains multiple predicted ARM repeats and a single C2 domain. Three-dimensional structures of previously characterized ARM repeats comprise three  $\alpha$  helices. For example, yeast importin- $\alpha$  contains a central region of 442 aa that contains 10 ARM repeats of 42 aa, forming a right-handed superhelix of helices that creates a surface for protein-protein interactions (18). By comparison, CSI1 has 10 predicted ARM repeats distributed unevenly across the entire protein (2,151 aa). We are not able to draw direct structural comparisons between CSI1 and other proteins containing ARM repeats.

## Methods

**Plant Materials and Growth Conditions.** *Arabidopsis thaliana* Columbia (Col-0) seeds were sterilized and germinated on Murashige and Skoog plates (1/2  $\times$  MS salts, 0.8% agar, 0.05% monohydrate 2-(N-Morpholino) ethanesulfonic acid, pH 5.7). Seedlings were grown vertically on the agar at 22  $^\circ$ C in darkness for 3 d before imaging. For soil-grown plants, seedlings were germinated on MS plates containing 1% sucrose and then were transferred to pots in a greenhouse at 22  $^\circ$ C under 16-h light and 8-h dark.

**Yeast Two-Hybrid Assay.** The yeast two-hybrid screen was carried out by Hybrigenics. CSI1 was identified as a CESA6-interactive protein. To confirm the interaction between CSI1 and CESA6, we subcloned the prey fragment into pACTII (28). The catalytic domains of CESA1, CESA3, and CESA6 were cloned into pAS1-CYH2 (28) using primers indicated in Table S2. The resulting constructs were cotransformed into yeast strain Y190 as pairs (Fig. 1B). Transformants were selected on SC-Trp-Leu-His plates. Positive interactions were tested by their ability to grow on SC-Trp-Leu-His plates





**Fig. 5.** Polarized light analysis of *csi1* mutants. (A and B) Polarized-light micrographs of (A) wild-type and (B) *csi1-1*-mutant roots. Images are of longitudinal-tangential sections through cortex and epidermis. Intensity is proportional to retardance, and color represents optical axis (azimuth) of the crystalline elements, as shown by the color wheel. Note the similar intensity and color for the two genotypes. (C) Quantification of retardance and azimuth ( $90^\circ$  is transverse to the root's long axis). Top panels show means  $\pm$  SEM ( $n = 3$  roots); bottom panels show the average SD ( $n \sim 200$  measurements)  $\pm$  SEM for each of the roots. Note the larger SDs for *csi1-1* mutants in the elongation and mature zones.

supplemented with 100 mM 3-aminotriazole (Sigma) and for GUS activity using a filter assay.

**Coexpression Analysis.** A coexpression network for *CSI1* (Cluster 86) was obtained from AraGenNet at <http://aranet.mpimp-golm.mpg.de/aranet/AraGenNet> (29) and trimmed to facilitate readability.

**GUS Construct and Staining.** A genomic DNA fragment (1.4 kb) upstream from the ATG start codon of *CSI1* was cloned into pCambia1305 GUS-Plus (Table S2) using BamHI and NcoI. The construct was transformed into *Arabidopsis* using *Agrobacterium*-mediated transformation. Transgenic plants were selected on hygromycin, stained for GUS activity in 100 mM sodium phosphate (pH 7.0), 10 mM EDTA, 1 mM ferri cyanide, 1 mM ferrocyanide, and 1 mM 5-bromo-4-chloro-3-indolyl  $\beta$ -D-glucuronic acid at  $37^\circ\text{C}$ , cleared in 70% ethanol, and observed under a Leica MZ12.5 stereomicroscope (Leica DFC420 digital camera).

**Construction of Transgenic Lines.** A 35S promoter in pH7WGR2 and pH7RWG2 (30) was replaced with a 3-kb promoter of *CSI1* using Spe I and SacI to create construct pYG104 and pYG105, respectively. The full-length coding sequence of *CSI1* (accession # NM\_12781.4) synthesized by DNA 2.0 was introduced into pYG104 and pYG105 using Gateway LR Clonase II (Invitrogen). The verified constructs (pYG106 and pYG107) then were introduced into *csi1-6* by *Agrobacterium*-mediated transformation. Complemented lines were selected for further analysis.

**Cellulose Measurement.** Cellulose was measured from 4-d-old etiolated seedlings using the Updegraff method (31). Data were collected from five technical replicates for each tissue sample.

**Microscopy.** Anthers from 5-wk-old plants or hypocotyls of 4-d-old dark-grown seedlings were mounted on stubs under a pressure of 50 Pa in an Hitachi TM-1000 scanning electron microscope. Roots of 8-d-old light-grown seedlings were wet-mounted on slides and viewed and photographed with a Leica DM5000B epifluorescence microscope (JH Technology). ImageJ software (W. Rasband, National Institute of Health, Bethesda, MD) was used for measurement of hypocotyls and root length. Whole 5-d-old seedlings were viewed on MS plates using a Leica MZ12.5 stereomicroscope (Leica DFC420 digital camera).

**Polarized Light Analysis.** Roots were prepared for polarized light analysis as described previously (27). Briefly, root tips were cryofixed in liquid propane, freeze-substituted in ethanol, embedded in butyl-methyl-methacrylate, and sectioned at  $2\text{-}\mu\text{m}$  thickness. Sections were imaged on an Interphako polarized-light microscope (Zeiss) equipped with an LC Polscope quantification system (Cambridge Research Instruments) implementing the universal compensator (26). This instrument operates in circularly polarized light and generates two images. The intensity of each pixel is proportional to birefringent retardance in the first image and to the azimuth of the optical axis of the crystalline elements in the second image. For display (Fig. 5) the two images are superimposed with pixel intensity giving retardance and color giving azimuth. Measurements were taken from subcellular areas of cell wall in cortex and epidermis, as viewed in longitudinal sections. Approximately 20 sections per root and three roots per genotype were measured. For analysis, the root was divided into meristem, elongation, and mature zone based on cell length. Azimuth was defined with respect to the local midline (longitudinal axis) of the root.

To assess the statistical significance of the polarized light observations, circular statistical techniques were used (32). Specifically, microfibril orientation was expressed as a unit vector corresponding to the doubled azimuth angle; to obtain the mean orientation, this unit vector was averaged over

each observation zone, and the result was halved. The angular deviation was calculated as half the (nonnegative) angular distance between the unit vectors for an individual patch,  $a$ , and the mean,  $b$ :  $\arccos[\cos(a - b)]$ . Zone-wise mean angular deviations of microfibrils were compared between genotypes using a Mann–Whitney U test, similar to the approach used to compare sample angular deviations reported by Wallraff (32). The distribution of zone mean angular deviations was skewed right, but the distribution became approximately normal after log transformation. Therefore, logarithms of mean angular deviations also were compared using an independent sample  $t$  test. The results of the  $t$  test on log-transformed data and the  $u$  test were the same.

**Isolation of T-DNA Insertion Line.** The identification of *csi1* knockout lines from the SIGnAL (Salk Institute Genomic Analysis Laboratory; <http://signal.salk.edu/cgi-bin/tdnaexpress>) collection was based on a combination of database searches and PCR amplification of T-DNA flanking regions. For T-DNA lines identified from the SIGnAL collection, seeds were obtained from the Arabidopsis Biological Resource Center (Ohio State University; <http://www.biosci.ohio-state.edu/~plantbio/Facilities/abrc/abrchome.htm>). PCR reactions were carried out to identify single plants for the T-DNA insertion. Primers used for T-DNA genotyping of *csi1* alleles are listed in Table S2.

**Confocal Microscopy and Image Analysis.** For analyses of microtubule dynamics, seeds were germinated on MS agar plates and grown vertically in darkness for 3 d at 22 °C. Seedlings were mounted between two coverslips in water. Imaging was performed on a Yokogawa CSUX1 spinning disk system featuring the DMI6000 Leica motorized microscope (13) and a Leica 100 $\times$ /1.4 NA oil objective. YFP was excited at 488 nm, and a band-pass filter (520/50 nm) was used for emission filtering. Image analysis was performed using MetaMorph (Molecular Devices) and Imaris (Bitplane) software.

For Imaris analysis, the contrast was enhanced and normalized for each slice within a movie using ImageJ. The enhanced movie was processed in Imaris 6.2.1 from Bitplane. Automated particle detection was performed to find particles with a diameter of  $\sim$ 230 nm, and tracks were generated over the lifetime of the particle. To filter noise in particle detection, only particles detected for 14 s (seven frames) were analyzed. The data for the total displacement and duration for each track in a movie were exported. Directional bias was analyzed by summing the vectors of all particles to determine the direction of greatest particle flux. The dot product of each particle's trajectory against this direction was calculated, and the velocity of each particle was plotted against its dot product. Greater differences between particle velocities going with and against the major axis result in a larger slope in a linear regression of this plot.

**Database Search and Sequence Alignment.** The predicted amino acid sequence of CSI1 (At2g22125) was retrieved from the Arabidopsis Information Resource (TAIR) database ([www.arabidopsis.org](http://www.arabidopsis.org)). This protein sequence was used to identify full-length CSI1-like proteins in the National Center for Biotechnology Information GenBank protein database using BLASTP ([www.ncbi.nlm.nih.gov/BLAST](http://www.ncbi.nlm.nih.gov/BLAST)). CSI1-like proteins (Table S3) were aligned using ClustalW implemented in MegAlign (DNASTAR); protein alignments then were used to generate the phylogenetic tree of CSI1-like proteins (MegAlign; DNASTAR).

**ACKNOWLEDGMENTS.** We thank H. Höfte (INRA, Versailles, France) for providing GFPESA3 line and S. Li and K. Hematy for helpful discussions. This work was supported in part by an award from The Balzan Foundation, Grants DE-FG02-03ER15421 (to T.I.B.) and DOE-FG02-03ER20133 (to C.R.S.) from the US Department of Energy, and the Energy Biosciences Institute. M.B., A.S., and S.P. were funded through the Max-Planck Gesellschaft.

- Brown RM, Jr, Willison JH, Richardson CL (1976) Cellulose biosynthesis in *Acetobacter xylinum*: Visualization of the site of synthesis and direct measurement of the in vivo process. *Proc Natl Acad Sci USA* 73:4565–4569.
- Mueller SC, Brown RM, Jr, Scott TK (1976) Cellulosic microfibrils: Nascent stages of synthesis in a higher plant cell. *Science* 194:949–951.
- Giddings TH, Jr, Brower DL, Staehelin LA (1980) Visualization of particle complexes in the plasma membrane of *Micrasterias denticulata* associated with the formation of cellulose fibrils in primary and secondary cell walls. *J Cell Biol* 84:327–339.
- Mueller SC, Brown RM, Jr (1980) Evidence for an intramembrane component associated with a cellulose microfibril-synthesizing complex in higher plants. *J Cell Biol* 84:315–326.
- Somerville C (2006) Cellulose synthesis in higher plants. *Annu Rev Cell Dev Biol* 22: 53–78.
- Persson S, et al. (2007) Genetic evidence for three unique components in primary cell-wall cellulose synthase complexes in Arabidopsis. *Proc Natl Acad Sci USA* 104: 15566–15571.
- Desprez T, et al. (2007) Organization of cellulose synthase complexes involved in primary cell wall synthesis in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 104: 15572–15577.
- Fagard M, et al. (2000) PROCUSTE1 encodes a cellulose synthase required for normal cell elongation specifically in roots and dark-grown hypocotyls of Arabidopsis. *Plant Cell* 12:2409–2424.
- Ellis C, Karafyllidis I, Wasternack C, Turner JG (2002) The Arabidopsis mutant *cev1* links cell wall signaling to jasmonate and ethylene responses. *Plant Cell* 14:1557–1566.
- Arioli T, et al. (1998) Molecular analysis of cellulose biosynthesis in Arabidopsis. *Science* 279:717–720.
- Taylor NG, Laurie S, Turner SR (2000) Multiple cellulose synthase catalytic subunits are required for cellulose synthesis in Arabidopsis. *Plant Cell* 12:2529–2540.
- Scheible WR, Eshed R, Richmond T, Delmer D, Somerville C (2001) Modifications of cellulose synthase confer resistance to isoxaben and thiazolidinone herbicides in Arabidopsis *lxr1* mutants. *Proc Natl Acad Sci USA* 98:10079–10084.
- Paredes AR, Somerville CR, Ehrhardt DW (2006) Visualization of cellulose synthase demonstrates functional association with microtubules. *Science* 312:1491–1495.
- Nicol F, et al. (1998) A plasma membrane-bound putative endo-1,4- $\beta$ -D-glucanase is required for normal wall assembly and cell elongation in Arabidopsis. *EMBO J* 17: 5563–5576.
- Persson S, Wei H, Milne J, Page GP, Somerville CR (2005) Identification of genes required for cellulose synthesis by regression analysis of public microarray data sets. *Proc Natl Acad Sci USA* 102:8633–8638.
- Nüsslein-Volhard C, Wieschaus E (1980) Mutations affecting segment number and polarity in *Drosophila*. *Nature* 287:795–801.
- Huber AH, Nelson WJ, Weis WI (1997) Three-dimensional structure of the armadillo repeat region of  $\beta$ -catenin. *Cell* 90:871–882.
- Conti E, Uy M, Leighton L, Blobel G, Kuriyan J (1998) Crystallographic analysis of the recognition of a nuclear localization signal by the nuclear import factor karyopherin alpha. *Cell* 94:193–204.
- Hatzfeld M (1999) The armadillo family of structural proteins. *Int Rev Cytol* 186: 179–224.
- Davletov BA, Südhof TC (1993) A single C2 domain from synaptotagmin I is sufficient for high affinity  $\text{Ca}^{2+}$ /phospholipid binding. *J Biol Chem* 268:26386–26390.
- Ochoa WF, et al. (2001) Structure of the C2 domain from novel protein kinase Cepsilon. A membrane binding model for  $\text{Ca}^{2+}$ -independent C2 domains. *J Mol Biol* 311:837–849.
- Benes CH, et al. (2005) The C2 domain of PKCdelta is a phosphotyrosine binding domain. *Cell* 121:271–280.
- Alonso JM, et al. (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* 301:653–657.
- Schindelman G, et al. (2001) COBRA encodes a putative GPI-anchored protein, which is polarly localized and necessary for oriented cell expansion in Arabidopsis. *Genes Dev* 15:1115–1127.
- Zhong R, Kays SJ, Schroeder BP, Ye ZH (2002) Mutation of a chitinase-like gene causes ectopic deposition of lignin, aberrant cell shapes, and overproduction of ethylene. *Plant Cell* 14:165–179.
- Oldenbourg R, Mei G (1995) New polarized light microscope with precision universal compensator. *J Microsc* 180:140–147.
- Baskin TI, Beemster GT, Judy-March JE, Marga F (2004) Disorganization of cortical microtubules stimulates tangential expansion and reduces the uniformity of cellulose microfibril alignment among cells in the root of Arabidopsis. *Plant Physiol* 135: 2279–2290.
- Kim J, Harter K, Theologis A (1997) Protein-protein interactions among the Aux/IAA proteins. *Proc Natl Acad Sci USA* 94:11786–11791.
- Mutwil M, et al. (2010) Assembly of an interactive correlation network for the Arabidopsis genome using a novel heuristic clustering algorithm. *Plant Physiol* 152: 29–43.
- Karimi M, Inzé D, Depicker A (2002) GATEWAY vectors for Agrobacterium-mediated plant transformation. *Trends Plant Sci* 7:193–195.
- Updegraff DM (1969) Semimicro determination of cellulose in biological materials. *Anal Biochem* 32:420–424.
- Batschelet E (1981) *Circular Statistics in Biology* (Academic, London), p 371.




PNAS




PNAS





**Fig. S3.** Morphology of *csi1* mutants. (A) RT-PCR analysis of *CSI1* mRNA expression in various transfer DNA (T-DNA) insertion lines in 1-wk-old seedlings. ACTIN2 was used as control. WT, Columbia-0. (B) Eight-day-old Col-0 (Top), *csi1-3*-mutant (Middle), and *csi1-6*-mutant (Bottom) seedlings were grown on 1/2 MS supplemented with 1% sucrose. (Scale bar: 5 mm.) (C) Quantification of primary root length in Col-0, *csi1-3*-mutant, and *csi1-6*-mutant seedlings. Measurement data were collected from about 50 seedlings in each genotype. (D and E) High magnification of primary root morphology in Col-0 (D) and *csi1-3*-mutant plants (E). (Scale bar: 200 μm.) (F) Quantification of primary root width in Col-0, *csi1-3*-mutant, and *csi1-6*-mutant seedlings. Measurement data were collected from about 50 seedlings in each genotype. (G) Adult plant morphology in *csi1* mutants: (Left to Right) Col-0 (wild-type) and *csi1-1*, *csi1-2*, *csi1-3*, *csi1-4*, and *csi1-6* mutants. (Scale bar: 5 cm.) Insets show enlarged image of siliques from wild-type (Col-0) and *csi1-3*-mutant plants. (Scale bar: 1 cm.) (H–K) Morphology of mature pollen grains in wild-type (H) and *csi1-3*-mutant (I) plants examined by SEM. (Scale bar: 200 μm.) Magnified SEM images of single pollen grains from wild-type (J) and *csi1-3*-mutant (K) plants. (Scale bar: 2 μm.)





**Fig. S4.** Both N- and C-terminal fusion constructs complement *csi1*-mutant phenotypes. (A) Schematic representation of N- and C-terminal fusion constructs. (B) Red fluorescent protein (RFP)-fusion constructs complemented *csi1*-mutant phenotype in dark-grown hypocotyls. (Scale bar: 5 mm.) (C) RFP-fusion constructs complemented *csi1*-mutant phenotype in adult plants: (Left to Right in B and C) wild-type (Col-0), *csi1-6* mutant, *CSI1::CSI1-RFP* in *csi1-6* mutant, and *CSI1::RFP-CSI1* in *csi1-6* mutant. (Scale bar: 5 cm.)

**Table S1. Reciprocal crosses indicate that *csi1* has pollen-specific transmission defect**


Allele	Pollen donor: +/- pollen recipient: +/+			Pollen donor: +/+ pollen recipient: +/-		
	<i>n</i>	+/+	+/-	<i>n</i>	+/+	+/-
Expected		50%	50%		50%	50%
<i>csi1-3</i>	86	82%	18%	62	51%	49%

**Table S2. DNA primers used in this study**

Transfer DNA line	Transfer DNA genotyping	RT-PCR analysis
<i>csi1-1</i> (SALK_136239)	F: CTAACAAAAGAGCAGTTGCCG R: CATCACACCCTATCGATTG	F: CTGAAAGTGGTGCCATAGAAGCTC R: CTTGTTAGACCACTGGAATTCT
<i>csi1-2</i> (SALK_122304)	F: TAGGCACACCAAAAGATGAAC R: GAGCTACAGAGCCTGCAACAC	F: CTGAAAGTGGTGCCATAGAAGCTC R: CTTGTTAGACCACTGGAATTCT
<i>csi1-3</i> (SALK_138584)	F: TAGGCACACCAAAAGATGAAC R: GAGCTACAGAGCCTGCAACAC	F: CTGAAAGTGGTGCCATAGAAGCTC R: CTTGTTAGACCACTGGAATTCT
<i>csi1-4</i> (SALK_047252)	F: GCTCCGAATTTTACCTGGAAG R: AGGAATTCAGCTTTCTCTGG	F: CTGAAAGTGGTGCCATAGAAGCTC R: CTTGTTAGACCACTGGAATTCT
<i>csi1-5</i> (SALK_051146)	F: CAATTTAGCCATCTGGTTGCTCTCGG R: CAACTAGTAAGGGAA-TGCCTTTTCGAG	F: GACTTTGCCGAGATCAGCCTATTG R: GTTCCGCGACTACCGTTACAGACA
<i>csi1-6</i> (SALK_115451)	F: CCGTCTTGGGTTCACTCTGCAAGGAGAA R: GGAAGCACTGCC-TTCACGTAGCATGTC	F: ATGACAAGTGCTCTTGATG R: CAATAGCATATTGGAAGCACTGCC
Y2H constructs	Forward primer	Reverse primer
CESA1	GGGATCCGTCAAGTTCCCAAATGGTAC	GTCGACTCAGACGATGGTGTGATATA
CESA3	CCATGGGAGATCAGTTTCCCAAGTGG	GTCGACTCAGATGGTGGTGTTCACATA
CESA6	CCATGGGACAGTTCCTAAATGGTAC	GTCGACTCAAACCACAGAGTTAATGTA
GUS construct	Forward primer	Reverse primer
<i>CSI1</i>	TTTAGGATCCCCATTGTTATGGGC (BamHI)	AATCTCCATGGAAGAGCACTTGTCAAATC (NcoI)

**Table S3. Accessions used in phylogenetic tree**


Name	Organism	Locus number	Accession number
AtCSI1	<i>Arabidopsis</i>	At2g22125	GI:79559926
AtCSI2	<i>Arabidopsis</i>	At1g44120	GI:15219352
AtCSI3	<i>Arabidopsis</i>	At1g77460	GI:186496022
PtCSIL1	<i>Populus trichocarpa</i>		GI:224083759
PtCSIL3	<i>Populus trichocarpa</i>		GI:224084912
PtCSIL2	<i>Populus trichocarpa</i>		GI:224096231
RcCSIL1	<i>Ricinus communis</i>		GI:255576178
RcCSIL2	<i>Ricinus communis</i>		GI:255558634
VvCSIL1	<i>Vitis vinifera</i>		GI:225438627
VvCSIL2	<i>Vitis vinifera</i>		GI:225459601
OsiCSIL1	<i>Oryza sativa Indica</i>		GI:218197824
OsiCSIL2	<i>Oryza sativa Indica</i>		GI:218185379
OsjCSIL1	<i>Oryza sativa Japonica</i>		GI:222635228
OsjCSIL2	<i>Oryza sativa Japonica</i>		GI:108864075
OsjCSIL3	<i>Oryza sativa Japonica</i>		GI:115484501
SbCSIL1	<i>Sorghum bicolor</i>		GI:242095194
SbCSIL2	<i>Sorghum bicolor</i>		GI:242067687
PpCSIL1	<i>Physcomitrella patens</i>		GI:168006338
PpCSIL2	<i>Physcomitrella patens</i>		GI:168024763



**Movie S1.** Time-lapse images of RFP-CSI1 at cell membrane in epidermal cells of dark-grown hypocotyls (quantified by velocity measurements in Fig. 3). Time series are 2 min long with images at 2-s intervals.

[Movie S1](#)





**Movie S2.** Time-lapse images of YFP-CESA6 at cell membrane in epidermal cells of dark grown *csi1-3* mutants (quantified by velocity measurements in Fig. 4). Time series are 2 min long with images at 2-s intervals.

[Movie S2](#)